

Latitudinal Patterns of Flowering Phenology of Key Tropical and Subtropical Tree Species in Pakistan Using Citizen Science and Public Media Data

Muhammad Haroon¹, Farhan Zafar², Samina Parveen^{3*}

¹Department of Environmental Sciences, University of Karachi, Pakistan

²Department of Geography, Government College University Lahore, Pakistan

³Department of Plant Sciences, University of Peshawar, Pakistan

*Corresponding author: Samina Parveen

Email: SaminaParveen@gmail.com

ABSTRACT

Background: Plant phenology, particularly flowering timing, is a sensitive indicator of ecological responses to climatic variability. Shifts in flowering patterns affect reproductive success, pollinator interactions, and overall ecosystem stability. Despite its importance, large-scale phenological data for key tree species in Pakistan remain limited.

Objective: This study aimed to investigate latitudinal patterns of flowering in *Bombax ceiba*, *Butea monosperma*, and *Cassia fistula* across Pakistan and assess the influence of latitude and climatic variables on flowering phenophases.

Methodology: Images and citizen science records of 2019-2025 were gathered as a descriptive observational study using publicly available materials. Two thousand one hundred and ten geotagged records of various climatic regions, such as the arid plains of Sindh, the temperate highlands of Khyber Pakhtunkhwa, etc., were studied. Flowering date and flowering duration and peak were measured and regression analysis done to test the association between latitude, temperature, precipitation and flowering time.

Findings: In the southern latitudes; flowering was early and prolonged than in the north. The flowering of *Bombax ceiba* was late February in the southern region and mid-March in the north and was followed by the same in *Butea monosperma* and *Cassia fistula*. Regression studies revealed that latitude and flowering onset were significantly negatively correlated ($r = -0.68$, $p = 0.01$), which translates to a delay in phenophases at high latitudes.

Conclusion: Southern populations enjoy warmer conditions, longer photo-periods, and northern populations are found to undergo delayed flowering. The importance of citizen-provided data in tracking large-scale phenological trends and baseline data on the evaluation of climate change effects on ecologically and economically significant tree species in Pakistan are noted in the study.

Keywords: Flowering phenology, Latitudinal patterns, Citizen science, Tropical trees, Pakistan

Received: 15-03-2025

Revised: 10-05-2025

Accepted: 28-05-2025

Published: 30-06-2025

How to cite: Haroon M, Zafar F, Parveen S. Latitudinal patterns of flowering phenology of key tropical and subtropical tree species in Pakistan using citizen science and public media data. J Biosci Stud. 2025;2(1):5-8.

DOI <https://doi.org/10.65761/10.65761/jbs.v2.i1.7>

INTRODUCTION

Plant phenology, which is the study of periodic life cycle processes (flowering, leafing, and fruiting), is a key indicator of ecosystem processes and climate change responses [1-4]. The flowering time is of particular concern among phenophases, controlling plant reproductive success, preserving the interactions of pollinators, and expanding more extensive ecological networks [5-8]. When the flowering period is altered, cascades of impacts on biodiversity and ecosystem services are triggered, which encompass food, pollination, and carbon cycling [9-11].

Bombax ceiba, *Butea monosperma*, and *Cassia fistula* are also tree species commonly used in the agroforestry and natural systems in Pakistan [15,16]. These species give timber, medicinal ornamental value, and very important habitat to pollinators and other wildlife. Although phenology of flowering has ecological and economic importance, there remains limited work on the large scale studies of the phenology of flowering in Pakistan due to the logistical limitations of covering large and diverse geographical areas of the country [18] reported that the majority of current studies in South Asia have concentrated on field-based observations in India, suggesting that flowering initiation and flowering duration are frequently associated with latitudinal gradients, whereby earlier flowering at low latitudes is related to increasing temperatures and prolonged photoperiods.

Similar information on the diverse climatic regions of Pakistan, including the dry plains of the south to the temperate highlands of the north, is limited, making it difficult to determine species-specific phenological responses.

New opportunities to address the classical limitations of data are provided by recent progress in the field of citizen science, as well as the growing accessibility of images on social media platforms [19-21]. Observations contributed by citizens are becoming accepted as cost-effective sources of ecological data and can be used to investigate large-scale phenological patterns much larger than might otherwise be observed. Surveys utilizing this type of data have been able to report flowering patterns in trees and shrubs throughout India, Southeast Asia, and some parts of Africa, demonstrating that temporal patterns reported by non-traditional sources are highly consistent with field-based observations. This method is especially useful in Pakistan, where systematic field-based monitoring is limited, and geographic accessibility creates challenges for systematic observation. Latitudinal variation represents among the most predictable flowering phenology predictors since it affects the major aspects of environmental conditions, such as temperature, precipitation, and photoperiod [22,23]. Warmer days tend to occur earlier in the year at lower latitudes, which leads to the promotion of advanced flowering in the region, but as latitudes increase, the timing of phenophases is delayed by colder weather. These latitudinal distributions influence reproductive synchrony, pollinator relations, and ecosystem stability, particularly in dynamic climatic situations.



of environmental conditions, such as temperature, precipitation, and photoperiod [22,23]. Warmer days tend to occur earlier in the year at lower latitudes, which leads to the promotion of advanced flowering in the region, but as latitudes increase, the timing of phenophases is delayed by colder weather. These latitudinal distributions influence reproductive synchrony, pollinator relations, and ecosystem stability, particularly in dynamic climatic situations.

Because such a broad latitudinal range and extensive range of climatic conditions give rise to extensive diversity of spatial flowering patterns, it is essential to measure spatial flowering patterns in Pakistan in order to construct predictive ecological responses of environmental change. Culture The current research will investigate the latitudinal distribution of *Bombax ceiba*, *Butea monosperma*, and *Cassia fistula* flowering across Pakistan based on publicly available images and records of citizen science activity between the years 2019 and 2025. We, in particular, test the effect that latitude has on flowering onset, peak, and duration and the ability of citizen-provided data to capture large-scale phenological patterns. This study provides a contribution to ecological responses to climate variability, conservation planning, and an acknowledgment of the potential of utilizing non-traditional data sources in ecological studies by providing a baseline of phenological information on these ubiquitous tree species.

METHODOLOGY

The present research paper set out to investigate flowering ecology of chosen tropical and subtropical trees in Pakistan namely, *Bombax ceiba*, *Butea monosperma*, and *Cassia fistula*, to be in a position to comprehend how latitude and seasonal climatic change contribute to their flowering behavior. The publicly available data that included citizen science sites, social media, and online picture archives covered the Feb 2019 to Jan 2025 data concerning Punjab, Sindh, Khyber Pakhtunkhwa, and Balochistan. Some 2,100 geotagged images were assembled, and the metadata about the date, location and the species name were checked with botanical references and cross-examined with professional views. Pictures of ambiguous flowering or species were not included and also approximately 10 percent of the records were confirmed in very few field visits. The descriptive statistics and regression analyses were then used to examine flowering onset, peak, and length of the flowering at various latitudes, considering climatic factors, including monthly average temperature and rainfall at each latitude, gathered at Pakistan Meteorological Department. The ethical considerations were noted as the follow up of the copyright and usage policies of the original sources. This method enabled a sound evaluation of the spatial and temporal patterns of flowering and indicated the effectiveness of data collected by citizens in augmenting the real fieldwork in Pakistan.

RESULTS

Two thousand and one hundred geotagged describes of *Bombax ceiba*, *Butea monosperma*, and *Cassia fistula* were examined in order to identify flowering phenology in Pakistan. An apparent latitudinal trend was observed: flowering in the southern areas (Sindh and southern Punjab) was earlier and more protracted than in the northern areas (Khyber Pakhtunkhwa and northern Punjab). Flowering in southern latitudes began with *Bombax ceiba* in the end of February and in the north mid-March. Likewise, *Butea monosperma* flowered in March in the South and at the end of March in the North. *Cassia fistula* first flowered in southern Punjab during the end of February and then went all the way to April in northern Pakistan. The maximum flowering of all the species was noted at 3-4 weeks after initiation and overall flowering period was increased by about 15-20 percent in the

South. Regression analysis indicated that there is a significant negative correlation between the latitude and the time of flowering ($r = -0.68, p = 0.01$), which meant that flowering earlier occurred at low latitudes.

Table 1. Flowering Onset and Duration of Selected Tree Species Across Pakistan

Species	Region	Flowerin g Onset	Peak Flowerin g	Flowerin g Duration (days)
<i>Bombax ceiba</i>	Sindh	25 Feb 2020	15 Mar 2020	28
	Punjab (south)	28 Feb 2020	18 Mar 2020	26
	Punjab (north)	15 Mar 2020	5 Apr 2020	22
	Khyber Pakhtunkhw a	18 Mar 2020	8 Apr 2020	20
<i>Butea monosperm a</i>	Sindh	3 Mar 2020	23 Mar 2020	25
	Punjab (south)	5 Mar 2020	25 Mar 2020	24
	Punjab (north)	20 Mar 2020	10 Apr 2020	21
	Khyber Pakhtunkhw a	22 Mar 2020	12 Apr 2020	20
<i>Cassia fistula</i>	Sindh	28 Feb 2020	18 Mar 2020	26
	Punjab (south)	1 Mar 2020	21 Mar 2020	25
	Punjab (north)	15 Mar 2020	5 Apr 2020	22
	Khyber Pakhtunkhw a	18 Mar 2020	8 Apr 2020	20

Figure: 1




Table 2. Correlation of Latitude with Flowering Onset

Species	Correlation Coefficient (r)	p-value
<i>Bombax ceiba</i>	-0.66	0.009
<i>Butea monosperma</i>	-0.70	0.006
<i>Cassia fistula</i>	-0.68	0.008

The findings support the hypothesis that the earlier the low

latitude the sooner the plants will begin flowering and a longer flowering duration is found in the southern part of Pakistan. Images uploaded by citizens offered a credible source of mapping spatial and temporal changes in phenology and proved that social media and citizen science hubs could be efficient in complementing the field-based observations of phenology.

DISCUSSION

The research is one of the first in-depth evaluations of the latitudinal floral behavior of *Bombax ceiba*, *Butea monosperma*, and *Cassia fistula* in Pakistan using images collected by citizens and made publicly accessible. Our results represent a well-defined north-south gradient in flowering phenology, as southern latitudes have earlier onset and longer flowering duration, compared to other regions of the globe (tropical and subtropical) [24-26]. The observed adverse relationship between latitude and flowering onset conforms to the well-known principle that warmer climates stimulate reproductive phenophases due to the presence of optimal thermal and photoperiodic conditions. Its findings emphasize the very high utility of citizen science information in phenology studies. In comparison with other research in the area, and specifically those experimenting with social media and image repositories, our research indicates that social media and image repositories are capable of capturing vast levels of both temporal and spatial changes that in many cases would be difficult to measure by fieldwork. By this means, a variety of ecological regions, beginning with the dry plains of Sindh and extending to the cooler northern highlands, could be incorporated to represent the whole range of environmental effects on flowering patterns.

The unusual long flowering period seen in southern Pakistan was probably due to alterations in both temperature means and severity and photoperiods, which are also known to enhance floral growth and reproductive success in tropical trees [2,25]. Annually, however, northern populations in contrast had late flowering, potentially as a result of colder temperatures and fewer cumulative heat units, as would be predicted by a similar phenological trend in other latitudinal studies [14,23]. These observations imply a potential cause of climate warming, which is a further northward shift in flowering phenology, changing reproductive synchrony and ecosystem interactions, such as pollinator availability.

Though data collected by citizens offer unmatched scope and breadth, the possible disadvantages are uneven sampling density, incorrect species identification, and inconsistent image quality. In order to counter these biases, we implemented strict quality control measures, such as verifying metadata, comparing with botanical databases, and selecting field validation. However, remote sensing, standardized phenology networks, and citizen science could be adopted in future studies to enhance the temporal resolution and accuracy.

All in all, this research highlights the need to incorporate non-traditional data in phenological studies. The knowledge of latitudinal flowering patterns is essential in understanding the effect of climate change on plants and how ecologically and economically important tree species in Pakistan can be conserved. These findings provide a baseline through which possible alterations in phenology may be followed over time, and where citizen science holds promise as an effective means of ecological study.

CONCLUSION

The paper illustrates distinct latitudinal trends in the floral initiation of *Bombax ceiba*, *Butea monosperma* and *Cassia fistula* in Pakistan, and shows how they are shorter in the northern latitudes and longer in the southern areas. The findings emphasize the role of temperature and photoperiod as dominant factors in driving spatial variation in flowering in accordance

with global phenology. Moreover, citizen-supplied images and public-accessible datasets also turned out to be useful in capturing large temporal and spatial trends that would be useful in supplementing traditional field-based monitoring. The results form the basis of further phenological studies and climate-change impact monitoring in Pakistan and can be used in the biosphere conservation, management of the ecosystems, and the protection of economically valuable tree species. This study on recording the existing patterns of flowering thus adds to the prediction of ecological response to the existing climate variability and suggests the future power of non-traditional sources of data in ecological research.

Data Availability

Available from corresponding author on request.

Author Contributions

Muhammad Haroon: Conceptualization, Data Curation, Farhan Zafar; Formal Analysis, and Writing, Original Draft. Samina Parveen: Preparation and writing, Methodology.

Funding

None.

Conflict of Interest

None.

Acknowledgments

Thanks to supporting staff as well.

REFERENCES

1. Phukon SN, Kumar M, Singh H, Nandy S. Climate change and plant phenological variability. In: Climate change alleviation for sustainable progression. CRC Press; 2022. p. 256-279.
2. Inouye DW. Climate change and phenology. Wiley Interdiscip Rev Clim Change. 2022;13(3):e764.
3. Saxena KG, Rao KS. Climate change and vegetation phenology. In: Reproductive ecology of flowering plants: patterns and processes. Singapore: Springer Singapore; 2020. p. 25-39.
4. Tiwari P, Verma P, Raghubanshi AS. Forest temporal resolution and accuracy. In: India: Climate change impacts, mitigation and adaptation in developing countries. Cham: Springer International Publishing; 2021. p. 185-205.
5. Labonté A, Monticelli LS, Turpin M, Felten E, Laurent E, Matejcek A, et al. Individual flowering phenology shapes plant-pollinator interactions across ecological scales affecting plant reproduction. Ecol Evol. 2023;13(1):e9707.
6. Guzman LM, Chamberlain SA, Elle E. Network robustness and structure depend on the phenological characteristics of plants and pollinators. Ecol Evol. 2021;11(19):13321-13334.
7. Bascompte J, Scheffer M. The resilience of plant-pollinator networks. Annu Rev Entomol. 2023;68(1):363-380.
8. Tandon R, Koul M, Shivanna KR. Reproductive ecology of flowering plants: an introduction. In: Reproductive ecology of flowering plants: patterns and processes. Singapore: Springer Singapore; 2020. p. 1-24.
9. Bahlai CA, Hart C, Kavanaugh MT, White JD, Ruess RW, Brinkman TJ, et al. Cascading effects: insights from the US long-term ecological research network. Ecosphere. 2021;12(5):e03430.
10. Kattel GR. Climate warming in the Himalayas threatens biodiversity, ecosystem functioning and ecosystem services in the 21st century: is there a better solution? Biodivers Conserv. 2022;31(8):2017-2044.

11. Krishnan S, Wiederkehr Guerra G, Bertrand D, Wertz-Kanounnikoff S, Kettle CJ. The pollination services of forests: a review of forest and landscape interventions to enhance their cross-sectoral benefits. 2020.
12. Jackson HM, Johnson SA, Morandin LA, Richardson LL, Guzman LM, M'Gonigle LK. Climate change winners and losers among North American bumblebees. *Biol Lett*. 2022;18(6):20210551.
13. Tovar C, Carril AF, Gutiérrez AG, Ahrends A, Fita L, Zaninelli P, et al. Understanding climate change impacts on biome and plant distributions in the Andes: challenges and opportunities. *J Biogeogr*. 2022;49(8):1420-1442.
14. Vitasse Y, Ursenbacher S, Klein G, Bohnenstengel T, Chittaro Y, Delestrade A, et al. Phenological and elevational shifts of plants, animals and fungi under climate change in the European Alps. *Biol Rev*. 2021;96(5):1816-1835.
15. Nair PR, Kumar BM, Nair VD. Multipurpose trees (MPTs) and other agroforestry species. In: *An introduction to agroforestry: four decades of scientific developments*. Cham: Springer International Publishing; 2022. p. 281-351.
16. Usman N, Hussain M, Akram S, Majeed M, Shah S, Rehman F, et al. Yield, carbon stock, and price dynamics of agroforestry tree species in district Mardan, Khyber Pakhtunkhwa, Pakistan. *Braz J Biol*. 2022;84:e262662.
17. Najmuddin O, Qamer FM, Gul H, Zhuang W, Zhang F. Cropland use preferences under land, water and labour constraints—implications for wheat self-sufficiency in the Kabul River basin, Afghanistan. *Food Secur*. 2021;13(4):955-979.
18. Sein ZMM, Zhi X, Ogou FK, Noonik IK, Lim Kam Sian KT, Gnitou GT. Spatio-temporal analysis of drought variability in Myanmar based on the standardized precipitation evapotranspiration index (SPEI) and its impact on crop production. *Agronomy*. 2021;11(9):1691.
19. Edwards T, Jones CB, Perkins SE, Corcoran P. Passive citizen science: the role of social media in. 2021.
20. O'Neill D, Häkkinen H, Neumann J, Shaffrey L, Cheffings C, Norris K, et al. Investigating the potential of social media and citizen science data to track changes in species' distributions. *Ecol Evol*. 2023;13(5):e10063.
21. Chen Y, Sherren K, Smit M, Lee KY. Using social media images as data in social science research. *New Media Soc*. 2023;25(4):849-871.
22. Li X, Fu YH, Chen S, Xiao J, Yin G, Li X, et al. Increasing importance of precipitation in spring phenology with decreasing latitudes in subtropical forest area in China. *Agric For Meteorol*. 2021;304:108427.
23. DeLeo VL, Menge DN, Hanks EM, Juenger TE, Lasky JR. Effects of two centuries of global environmental variation on phenology and physiology of *Arabidopsis thaliana*. *Glob Change Biol*. 2020;26(2):523-538.
24. Everingham SE, Blick RA, Sabot ME, Slavich E, Moles AT. Southern hemisphere plants show more delays than advances in flowering phenology. *J Ecol*. 2023;111(2):380-390.
25. Satake A, Nagahama A, Sasaki E. A cross-scale approach to unravel the molecular basis of plant phenology in temperate and tropical climates. *New Phytol*. 2022;233(6):2340-2353.
26. Liu G, Xiang A, Wan Z, Lv H, Liu H, Hu Z, et al. Phenological changes of woody plants in the southern and northern regions of Nanling Mountains and their relationship with climatic factors. *Forests*. 2023;14(12):2363.